



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C07K 14/785, A61P 11/00	A1	(11) International Publication Number: WO 00/47623 (43) International Publication Date: 17 August 2000 (17.08.00)
(21) International Application Number: PCT/EP00/01044 (22) International Filing Date: 9 February 2000 (09.02.00) (30) Priority Data: MI99A000275 12 February 1999 (12.02.99) IT (71) Applicant (for all designated States except US): CHIESI FARMACEUTICI S.P.A. [IT/IT]; Via Palermo, 26/A, I-43100 Parma (IT). (72) Inventors; and (75) Inventors/Applicants (for US only): CURSTEDT, Tore [SE/IT]; Via Palermo, 26/A, I-43100 Parma (IT). JOHANSSON, Jan [SE/IT]; Via Palermo, 26/A, I-43100 Parma (IT). JÖRNVALL, Hans [SE/IT]; Via Palermo, 26/A, I-43100 Parma (IT). ROBERTSON, Bengt [SE/IT]; Via Palermo, 26/A, I-43100 Parma (IT). VENTURA, Paolo [IT/IT]; Via Palermo, 26/A, I-43100 Parma (IT). (74) Agents: MINOJA, Fabrizio et al.; Bianchetti Bracco Minoja S.r.l., Via Rossini, 8, I-20122 Milan (IT).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: ARTIFICIAL PEPTIDES HAVING SURFACE ACTIVITY AND THE USE THEREOF IN THE PREPARATION OF ARTIFICIAL SURFACTANT (57) Abstract <p>SP-C analogue artificial peptides having surface activity are disclosed and their use for preparing artificial surfactants useful in the treatment of respiratory distress syndrome (RDS) and other diseases related to surfactant deficiency or dysfunction.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

**ARTIFICIAL PEPTIDES HAVING SURFACE ACTIVITY AND THE
USE THEREOF IN THE PREPARATION OF ARTIFICIAL
SURFACTANT**

The present invention provides new artificial peptides having surface activity. In particular, the invention provides SP-C analogues which, once combined with suitable lipids, are particularly effective in reducing surface tension at the air-liquid interface.

5 Thus, the peptides of the invention may be used in combination with lipids, and optionally in combination with SP-B or an active analogue thereof or a substitute of SP-B, for preparing artificial surfactants useful in the treatment of respiratory distress syndrome (RDS), other surfactant deficiencies or dysfunction,
10 related pulmonary diseases such as pneumonia, bronchitis, asthma, meconium aspiration syndrome and also other diseases such as serous otitis media (glue ear).

Background of the invention

Pulmonary surfactant reduces surface tension at the air-liquid
15 interface of the alveolar lining, preventing the lungs from collapsing at end expiration. Surfactant deficiency is a common disorder in premature infants and causes respiratory distress syndrome (RDS), which can be effectively treated with natural surfactants extracted from animal lungs (Fujiwara, T. and
20 Robertson B. (1992) *In*: Robertson, B., van Golde, L.M.G. and Batenburg, B. (eds) Pulmonary Surfactant: From Molecular Biology to Clinical Practice Amsterdam, Elsevier, pp. 561-592). The main constituents of these surfactant preparations are phospholipids such as 1,2-dipalmitoyl-*sn*-glycero-3-phosphocholine (DPPC),

phosphatidylglycerol (PG) and the hydrophobic surfactant proteins B and C (SP-B and SP-C). The hydrophilic surfactant proteins SP-A and SP-D which are C-type (Ca^{2+} -dependent) collagenous lectins and thought to act primarily in the host-defence system, are normally not included in the surfactant preparations due to the organic solvent extraction procedures employed.

SP-B and SP-C constitute only about 1-2 % of the surfactant mass, but are still able to exercise dramatic improvements on surface activity, compared to pure lipid preparations (Curstedt, T. et al. (1987) *Eur. J. Biochem.* **168**, 255-262; Takahashi, A., Nemoto, T. and Fujiwara, T. (1994) *Acta Paediatr. Jap.* **36**, 613-618). The primary and secondary structures of SP-B and SP-C and a tertiary structure of SP-C in solution have been determined (see 4). SP-B is composed of two identical polypeptide chains of 79 amino acids, connected with an interchain disulphide bridge (Curstedt, T. et al. (1990) *Proc. Natl. Acad. Sci. U.S.A.* **87**, 2985-2989; Johansson, J., Curstedt, T. and Jörnvall, H. (1991) *Biochemistry* **30**, 6917-6921). Each monomeric chain has three intrachain disulphide bridges and at least four amphipathic helices exhibiting one polar and one unpolar face through which SP-B may interact with two lipid bilayers and bring them into close proximity (Andersson, M. et al. (1995) *FEBS Lett.* **362**, 328-332). SP-C is a lipoprotein composed of 35 amino acid residues with an α -helical domain between residues 9-34 (Johansson, J. et al. (1994) *Biochemistry* **33**, 6015-6023). The helix is composed mostly of valyl-residues and is embedded in a lipid bilayer and oriented in parallel with the lipid acyl chains (Vandenbussche, et al. (1992)

Eur. J. Biochem. **203**, 201-209). Two palmitoyl groups are covalently linked to cysteine residues in positions 5 and 6 in the N-terminal part of the peptide (Curstedt, T. et al. (1990) Proc. Natl. Acad. Sci. U.S.A. **87**, 2985-2989). The two conserved positively charged residues, arginine and lysine, at positions 11 and 12, possibly interact with the negatively charged head groups of the lipid membrane, thus increasing its rigidity. The rigidity of the lipid-peptide interaction may be decreased towards the C-terminal end, since it contains small or hydrophobic residues only, making this part potentially more mobile in a phospholipid bilayer. SP-C is thought to influence the thickness and fluidity of the surrounding lipids *via* the extremely stable poly-valyl helix (Johansson, J. and Curstedt, T. (1997) Eur. J. Biochem. **244**, 675-693)

State of the art

Since surfactant preparations obtained from animal tissue present some drawbacks, like their availability in limited amounts and the possibilities that they contain infectious agents and induce immunological reactions, attempts have been made to create artificial surfactants (Johansson, J. and Curstedt, T. (1997) Eur. J. Biochem. **244**, 675-693; Johansson, J. et al. (1996) Acta Paediatr. **85**, 642-646), usually from synthetic lipids and hydrophobic proteins.

Previous work has demonstrated that synthetic SP-C may not fold like the native peptide into an α -helical conformation necessary for optimal surface activity (Johansson, J. et al. (1995) Biochem. J. **307**, 535-541), and therefore do not interact properly with the surfactant lipids.

Consequently, synthetic SP-C analogues do not fold like the

native peptide and do not interact properly with the surfactant lipids. To circumvent this problem, several attempts have been made to modify the sequence, for instance by replacing all helical Val residues in native SP-C with Leu, which strongly favour α -helical conformation. The corresponding transmembranous analogue, SP-C(Leu) showed good spreading at an air-liquid interface when combined with DPPC:PG:PA (68:22:9) (w/w/w). However, the maximum surface tension value during cyclic surface compression (γ_{\max}) was significantly higher than that of native surfactant. It was furthermore not possible to prepare lipid-peptide mixtures of higher concentrations than about 20 mg/ml, probably due to formation of peptide oligomers (Nilsson, G. et al. (1998) Eur. J. Biochem. 255, 116-124). Others have synthesised bioactive polyleucine SP-C analogues of different lengths (Takei, T. et al. (1996) Biol. Pharm. Bull. **19**, 1550-1555). In the latter studies neither self oligomerisation nor problems in producing samples of high lipid concentration were reported.

Different publications deal with the problem of providing peptidic analogues of natural surfactant peptides, giving a number of different solutions. Among those publications, WO93 21225, EP 733 645, WO96 17872, in the name of Tokyo Tanabe, disclose peptides analogues of natural SP-C, which in general differ from the native peptide regarding the sequence of the N-terminal part.

Scripps Research Institute patent applications WO89 06657 and WO92 22315 disclose SP-B analogues having alternating hydrophobic and hydrophilic amino acid residues. Among others, a peptide alternating Leucine and lysine residues (KL₄) is claimed.

Clercx A. et al., Eur. J. Biochem 229, 465-72, 1995, disclose

peptides of different lengths corresponding to the N-terminal of porcine SP-C and hybrid peptides derived from porcine SP-C and bacteriorhodopsin.


Johansson J. et al., Biochem. J. 307, 535:41, 1995, disclose synthetic peptides that differ from the native porcine SP-C by the substitution of some amino acids.

WO89/ 04326 in the name of California Biotechnology - Byk Gulden, and WO91/18015 in the name of California Biotechnology - Scios Nova, disclose SP-C analogues containing an initial N-terminal sequence in which the two Cys of natural SP-C are replaced by two Ser.

Description of the invention

It has now been found that SP-C analogue peptides which combine the following features: i) substitution of Val residues with other neutral and hydrophobic residues; ii) substitution of Cys residues with Ser residues; iii) replacement of some of the neutral amino acid residues with bulky or polar residues, show particularly favourable properties for surface tension reduction. In particular it has been found that the latter feature, in virtue of the positive charges conferred by the polar residues or the steric hindrance conferred by the bulky substituents, allow to avoid self-oligomerisation.

As follows, according to a first aspect, the invention provides SP-C analogues having the following general formula (I), using the one-letter amino acid code:



wherein:

X is an amino acid selected from the group consisting of

V, I, L, Nle (norleucine);

B is an amino acid selected from the group consisting of Ornithine, K, I, W, F, Y, Q, N;

5 Z is an amino acid selected from the group consisting of S, C, F where Ser or Cys residues are optionally linked via ester or thio-ester bonds with acyl groups containing 12-22 carbon atoms linked.

a is an integer from 1 to 19

b is an integer from 1 to 19

10 c is an integer from 1 to 21

d is an integer from 0 to 20

e is 0 or 1

f is 0 or 1

n is 0 or 1

15 m is 0 or 1

with the conditions:

- $n + m > 0$,

- $f \geq e$;

20 - $(X_a B)_n (X_b B)_n (X_c B)_m X_d$ is a sequence having a maximum of 22 amino acids, preferably from 10 to 22.

Preferred peptides of Formula (I) have the following sequences:

(Ia) FGIPSSPVHLKRX₄BX₄BX₄BXGALLMGL

(Ib) FGIPSSPVHLKRX₅BX₅BX₄GALLMGL

(Ic) FGIPSSPVHLKRX₄BX₁₁GALLMGL

25 (Id) FGIPSSPVHLKRX₈BX₇GALLMGL

(Ie) FGIPSSPVHLKRX₁₁BX₄GALLMGL

Among the sequences (Ia) - (Ie), those having B = Lys or Phe and X = Leu, Ile or Nle are preferred.

According to preferred embodiments, peptides of formula (Ia) - (If) have the following sequences, respectively:

FGIPSSPVHLKRLILKLLLLKILLKLGALLMGL [SP-C (LKS)]

FGIPSSPVHLKRLILLLKLLLLIKLLILGALLMGL [SP-C (LKS)₁]

5 FGIPSSPVHLKRLILKLLLLLILLILGALLMGL [SP-C (LKS)₂]

FGIPSSPVHLKRLILLLLLLKILLILGALLMGL [SP-C (LKS)₃]

FGIPSSPVHLKRLILLLLLLLIKLLILGALLMGL [SP-C (LKS)₄]

FGIPSSPVHLKRLILFLLLLFILLFLGALLMGL [SP-C (LFS)]

10 In a more preferred embodiment of the invention, the Ser residues are covalently linked with acyl groups containing 12-22 carbon atoms.

Peptides of formula (I) may be prepared by synthetic methods or recombinant techniques.

Conventional synthetic methods are described, for instance,
15 in Schroeder et al., "The peptides", vol. 1, Academic Press, 1965; Bodanszky et al., "Peptide synthesis", Interscience Publisher, 1996; Baramy & Merrifield, "The peptides; Analysis, Synthesis, Biology", vol. 2, chapter 1, Academic Press, 1980. Said techniques include peptide synthesis in solid phase, in solution, organic
20 chemistry synthetic methods, or any combination thereof.

S- or O- acylated peptides are preferably synthesized by treatment of the non-acylated peptides with acyl chloride in neat trifluoroacetic acid as described in Yousefi-Salakdeh et al. Biochem J 1999, 343, 557-562. After synthesis and purification,
25 the synthetic peptides were biochemically and biophysically characterised, as reported in the following section "Examples".

The activity of the peptides of the invention in reducing surface tension has been evaluated in combination with lipids and

phospholipids, SP-B, analogues of SP-B or substitutes of SP-B. In particular, the peptides have been combined with DPPC (1,2-dipalmitoyl-*sn*-glycero-3-phosphocholine)/PG (phosphatidylglycerol)/PA (palmitic acid) with or without SP-B, an active analogue thereof and polymyxins.

The results of pulsating-bubble surface activity tests clearly show that the synthetic peptides according to the present invention strongly decrease minimum and maximum surface tension during cyclic surface compression (γ_{\min} and γ_{\max}) to values comparable with those obtained using surfactants from natural sources.

The addition of SP-B or active analogue thereof to the mixture peptide/lipids-phospholipids gave particularly favourable results. Furthermore it has been surprisingly found that polymyxins, in particular polymyxin B, act as substituents of SP-B and their addition gave comparable results to those achieved with SP-B.

According to a second aspect, the invention provides a synthetic surfactant comprising one or more peptides of formula (I), in admixture with lipids and/or phospholipids and optionally SP-B, an active derivative thereof or polymyxins. Suitable lipids/phospholipids may be selected from the group consisting of phosphatidylcholines (preferably DPPC) PG, PA, triacylglycerols, sphingomyelin.

In an even more preferred embodiment of the invention, surfactant mixtures containing the peptide in which palmitoyl chains are O-covalently linked to the Ser residues should be used. It has been found that surfactant mixtures containing a dipalmitoylated form of the reference peptide (SP-C(Leu)) exhibit

higher surface film stability and increased size of the surface-associated lipid reservoir, compared to mixtures containing the corresponding non-palmitoylated peptide, as measured by captive bubble system. In the samples containing 5% of dipalmitoylated peptide, the γ_{\min} was below 1.5 mN/m and the films very stable, as the surface tension increased by less than 0.5 mN/m within 10 min at constant bubble volume. On the contrary, the γ_{\min} for the non-palmitoylated peptide was approx. 5 mN/m and the films were less stable as observed by frequent bubble clicking at low surface tensions. Moreover, after subphase depletion for samples that contain non-palmitoylated peptide, the ability to reach near zero stable surface tension was lost after a few adsorption steps, whereas with the dipalmitoylated peptide the film quality did not deteriorate even after more than 10 expansion steps and the incorporation of reservoir material equivalent to more than two monolayers. The improved surface activity of dipalmitoylated peptides was also demonstrated by pulsating bubble surfactometer. In addition, the presence of acyl groups was found to further reduce the tendency to form oligomers. This finding is very important, as during preparation of artificial surfactants, peptide oligomerisation has been found to hinder preparation of the mixtures at higher concentrations than 20 mg/ml (Nilsson *et al.* Eur J Biochem 1998, 255, 116-124).

The synthetic surfactant may be prepared by mixing solutions or suspensions of peptides and lipids and by subsequently drying the mixture.

At the occurrence, the dry mixture may be suspended, dispersed or administered as such to subjects in need of treatment

for surfactant deficiency.

The synthetic surfactant will be preferably administered endotracheally or via aerosol. The latter form of administration will require the combination of small particles of surfactant with
5 suitable inert propellant. Other forms of administration, like nebulization or spraying of stable solutions/suspensions of surfactant are also included within the scope of the invention.

According to a further aspect, the invention provides the use of the described peptides for the preparation of a surfactant agent
10 to be used in all cases of adult or neonatal surfactant deficiency or dysfunction, related pulmonary diseases such as pneumonia, bronchitis, asthma, meconium aspiration syndrome and also other diseases such as serous otitis media (glue ear).

Typically, the surfactant agent will be used, preferably upon
15 endotracheal administration, in the treatment of respiratory distress syndrome which frequently affects premature infants.

The following examples illustrate the invention in more details.

Example 1

20 **Peptide synthesis and purification**

An analogue of SP-C, SP-C(LKS) (Fig. 1) was synthesised by use of stepwise solid phase technology and the *tert*-butyloxycarbonyl chemistry (Kent, S.B.H. (1988) Annu. Rev. Biochem. **57**, 957-989) in an Applied Biosystems 430A instrument.
25 Cleavage of the resin-peptide bond and deprotection of the side-chains were carried out in anhydrous hydrogen fluoride/methoxybenzene/dimethylsulfide, 10:1:1 (v/v/v) for 1.5 h at 0°C. Protecting groups and scavengers were removed by

repeated extraction with diethyl ether and the peptide was subsequently extracted from the resin by dichloromethane/trifluoroacetic acid (TFA) 3:1 (v/v) followed by rotary evaporation. The crude peptide extract was redissolved at a concentration of 100 mg/ml in chloroform/methanol 1:1 (v/v) containing 5% H₂O. An aliquot of 10 mg was applied on a Sephadex LH-60 column (40 x 1 cm) in the same solvent (Curstedt, T. et al. (1987) Eur. J. Biochem. **168**, 255-262). Fractions of 2.5 ml were collected and absorbencies at 214 and 280 nm were measured. Identification and quantitation were performed by amino acid analysis.

For acylation, the purified peptide (typically about 5 mg) is dried, dissolved in distilled TFA (100 µl) and acyl chloride (10-20 equivalents compared to peptide) is added. After 10 minutes the reaction is quenched with 80% aqueous ethanol (1.9 ml). Purification of acyl peptides is performed using chromatography over Lipidex 5000 in ethylene chloride/methanol 1:4 (v/v) followed by reversed-phase HPLC over a C18 column using a linear gradient of 2-propanol/0.1% TFA running into 60% (aqueous) methanol/0.1% TFA or 75% (aqueous) ethanol/0.1% TFA.

Example 2

Biochemical characterisation

The purity of the peptide was checked by sodium-dodecylsulphate (SDS) polyacrylamide gel electrophoresis (PAGE) (Phast-system, Pharmacia, Sweden) and by reversed phase high performance liquid chromatography (HPLC), using a C₁₈ column and a linear gradient of 60 % aqueous methanol /0.1 % TFA and isopropanol/ 0.1 % TFA (Gustafsson, M. et al. (1997) Biochem. J.

326, 799-806).

Molecular masses were determined by matrix-assisted laser desorption ionisation-time-of-flight (MALDI-TOF) mass spectrometry (Lasermat 2000, Finnigan MAT) calibrated with
5 vasoactive intestinal peptide (M_r 3326.8).

Peptide secondary structure was investigated using circular dichroism (CD) spectroscopy (Jasco-720 Jasco, Japan). After solubilisation with trifluoroethanol (TFE) spectra were recorded from 260 to 184 nm with a scan speed of 20 nm/min and a
10 resolution of 2 data points/nm. The residual molar ellipticity was calculated and expressed in $\text{kdeg} \times \text{cm}^2/\text{dmol}$. Molar ellipticities at 208 and 222 nm were utilised for estimating the content of helical structure (Barrow, C.J. et al. (1992) *J. Mol. Biol.* **225**, 1075-1093).

Secondary structure investigations of SP-C(LKS) using CD
15 spectroscopy showed a spectrum typical for α -helical peptides and an α -helical content of approximately 75 % was estimated from the 208 nm and 222 nm minima. The secondary structure remained stable following stepwise dilution with H_2O until 12 % TFE provided that the peptide was solubilized in neat TFE.

20 SDS-PAGE of SP-C(LKS) showed a single band similar to native SP-C while SP-C(Leu) which lacks Lys in the helical part forms oligomers. In contrast to our experience with SP-C(Leu)/lipid mixtures, which are difficult to solubilize in higher concentrations than 20 mg/ml (Nilsson, G. et al. (1998) *Eur. J. Biochem.* **255**, 116-124.), it was possible to make a SP-C(LKS)/lipid mixture with a lipid concentration of 80 mg/ml and a
25 polypeptide/lipid ratio of 0.03.

Example 3

Preparation of peptide/lipid mixtures

DPPC, PG and PA were all purchased from Sigma Chemical Co. (St Louis, MO). The lipids, dissolved in chloroform/methanol
5 98:2 (v/v), were mixed in the proportions DPPC:PG:PA 68:22:9 (w/w/w) or DPPC/PG 7:3 (w/w). Surfactant preparations were prepared by adding SP-C(LKS) alone or SP-C(LKS) and SP-B, to each of the lipid mixtures, at total polypeptide/lipid weight ratios of 0-0.05. The mixtures were evaporated under nitrogen and
10 resuspended in 150 mmol/l NaCl or in 10 mmol/l Hepes buffer pH 6.9 containing 140 mmol/l NaCl and 2.0 mmol/l CaCl_2 , at lipid concentrations of 10 — 80 mg/ml. Repeated freezing and sonication (50 W, 48 kHz) were performed until homogeneous suspensions were achieved. In some cases the final suspensions
15 were incubated at 45°C for 1 h.

Surfactant preparations suspended in 150 mmol/l NaCl have a pH of 3.5-5.5. The lower pH-values 3.5-4.5 were observed in preparations containing SP-B. Since native SP-B is purified using acidified organic solvents (Curstedt, T. et al. (1987) Eur. J.
20 Biochem. **168**, 255-262) small amounts of acid may remain in the preparations. Near physiological pH was obtained by suspending the surfactant preparation in Hepes buffer pH 6.9, containing 140 mmol/l NaCl and 2 mmol/l CaCl_2 (Table 1). Compared to the corresponding preparations in unbuffered saline there were no
25 changes of γ_{\max} or γ_{\min} when DPPC/PG 7:3 (w/w) was used as the lipid mixture. However when PA was included in the lipid mixture both γ_{\max} and γ_{\min} increased at the higher pH (Tables 1 and 2).

Table 1: Surface properties of preparations of artificial surfactant in physiological saline solution

Measurements were carried out directly after the preparation of the samples or after incubation for 1 hour at 45° C. The phospholipids conc. was 10 mg/ml in NaCl 150 mmol/l. The recordings were obtained at different periods of time with a pulsating bubble surfactometer at 37° C, 50% surface compression and at a rate of 40 cycles per min. The values are the mean (standard deviation) of 3-5 measurements. Abbreviations are defined in the text.

Surfactant preparation			Surface tension (mN/m)						
SP-C (LKS) (% w/w)	SP-B (% w/w)	Phospholipids	Incubation temperature	7,5 s		1 min		5 min	
				γ_{\min}	γ_{\max}	γ_{\min}	γ_{\max}	γ_{\min}	γ_{\max}
3	-	DPPC/PG/PA	-	<1	41(1)	<1	41(1)	<1	41(0)
3	-	DPPC/PG/PA	45°C	<1	41(1)	<1	41(1)	<1	41(1)
3	2	DPPC/PG/PA	-	<1	33(2)	<1	33(2)	<1	33(2)
3	2	DPPC/PG/PA	45°C	<1	34(1)	<1	34(2)	<1	25(2)
3	-	DPPC/PG	-	12	39(5)	14(4)	42(4)	9(5)	42(2)
3	-	DPPC/PG	45°C	8(3)	35(5)	9(3)	39(5)	6(4)	42(3)
3	2	DPPC/PG	-	2(1)	31(1)	2(1)	31(3)	2(1)	33(1)
3	2	DPPC/PG	45°C	3(3)	29(3)	1(1)	33(2)	1(0)	36(1)
3	1	DPPC/PG	-	<1	24(4)	1(2)	26(4)	<1	31(1)
3	0.5	DPPC/PG	45°C	4(2)	29(1)	4(3)	29(2)	3(1)	34(2)

Table 2: Surface properties of preparations of artificial surfactant in buffered salt solution

Measurements were carried out on samples containing phospholipids at the conc. of 10 mg/ml in Hepes buffer (pH 6.9), in turn containing NaCl 140 mmol/l and CaCl₂ 2.0 mmol/l. The recordings were obtained at different periods of time with a pulsating bubble surfactometer at 37° C, 50% surface compression and at a rate of 40 cycles per min. The values are the mean (standard deviation) of 3-5 measurements. Abbreviations are defined in the text.

Surfactant preparation		Surface tension (mN/m)			
SP-C (LKS) (% w/w)	SP-B (% w/w)	Phospholipids	7,5 s ----- γ_{\min} γ_{\max}	1 min ----- γ_{\min} γ_{\max}	5 min ----- γ_{\min} γ_{\max}
3	-	DPPE/PG/PA	4(1) 44(2)	5(2) 47(2)	7(2) 50(1)
3	2	DPPE/PG/PA	3(3) 38(3)	4(4) 40(2)	3(3) 44(2)
3	-	DPPE/PG	15(3)	16(2) 42(3)	13(3) 44(3)
3	2	DPPE/PG	39(4) 26(3)	1(1) 29(3)	<1 35(1)

Example 4.

Preparation of phospholipid mixtures with SP-C(LKS) and polymyxin B.

5 DPPC and PG were purchased from Sigma Chemical Co (St Louis, MO). The phospholipids, dissolved in chloroform/methanol 98:2 (v/v), were mixed in the proportions DPPC/PG 7:3 (w/w). SP-C (LKS) was added to the phospholipid mixtures, at a total polypeptide/phospholipid weight ratio of 0.03. The mixtures were evaporated under nitrogen and resuspended at room temperature
10 in 10 mmol/l Hepes buffer pH 6,9 containing 140 mmol/l NaCl and 2.0 mmol/l CaCl_2 or in the same buffer containing 0.01% polymyxin B (PxB) (Sigma Chemical Co, St Louis, MO). Repeated freezing and sonication (50 W, 48 kHz) were performed until homogeneous suspensions were achieved. The final phospholipid
15 concentration for both preparations was 10 mg/ml. Addition of PxB decreased both γ_{\min} and γ_{\max} and optimal surface activity was obtained (Table 3).

Table 3. Surface properties of artificial surfactant with and without polymyxin B

The recordings were obtained at different periods of time with a pulsating bubble surfactometer at 37°C and 50% surface compression at a rate of 40 cycles per min. The values are the mean (SD) of 5-11 measurements. Abbreviations are defined in the text.

Surfactant preparation		Surface tension (mN/m)			
SP-C (LKS) (% w/w)	PxB (% w/w)	Phospholipids	7,5 s ----- γ_{\min} γ_{\max}	1 min ----- γ_{\min} γ_{\max}	5 min ----- γ_{\min} γ_{\max}
3	-	DPPC/PG	15(3) 3(2)	16(2) 2(2)	13(3) 1(1)
3	1	DPPC/PG	39(4) 29(3)	42(3) 31(4)	44(3) 34(2)

Example 5

Biophysical characterisation

Surface spreading kinetics were measured at about 34-37 °C with a Wilhelmy surface balance (Biegler, Vienna, Austria). Surface tension was monitored for 10 min using a platinum plate connected to a strain gauge and inserted 1 mm in to a hypophase of 20 ml of 150 mmol/l NaCl in a teflon trough. The suspensions were added as droplets, totally 1 mg of lipids, onto the hypophase, 4 cm from the platinum plate.

Kinetic measurements of 3 weight % SP-C(LKS) in DPPC/PG, 7:3 (w/w), using the Wilhelmy balance showed a rapid spreading with a surface tension of 28 mN/m after 3 s (Fig. 2). The spreading was somewhat slower using 1 weight % SP-C(LKS) in the same lipid mixture (data not shown). Addition of 2 weight % SP-B did not significantly change spreading velocity or equilibrium surface tension (Fig. 2). No improvements were observed after incubation of the mixture for 1 h at 45°C (data not shown). Similar results were obtained with DPPC:PG:PA, 68:22:9 (w/w/w) as the lipid mixture (data not shown).

Dynamic surface tension was recorded using a pulsating bubble surfactometer (Surfactometer International, Toronto, Canada) at 37°C during 50 % cyclic compression of the bubble surface and at a frequency of 40 cycles per min. All measurements were performed for 5 min and at a lipid concentration of 10 mg/ml. The pressure gradients across the bubble wall were measured at specific time intervals were measured and used to calculate surface tensions at minimum (γ_{\min}) and maximum (γ_{\max}) bubble size.

In the pulsating bubble surfactometer 3 weight % SP-C(LKS) in DPPC:PG:PA, 68:22:9 (w/w/w,) produced a surface tension of less than 1 mN/m at minimum bubble radius (γ_{\min}) while a γ_{\min} of 9-14 mN/m was observed with 3 weight % SP-C(LKS) in DPPC:PG, 7:3 (w/w) (Table 1). The surface tension at maximum bubble radius (γ_{\max}) was about 40 mN/m in both cases. Addition of 2 weight % SP-B gave γ_{\max} -values of 31-33 mN/m and γ_{\min} of 0-2 mN/m for both lipid preparations. These values are very similar to those obtained with surfactant preparations isolated from natural sources (Robertson, B. et al. (1990) Prog. Respir. Res. **25**, 237-246). Incubation of the preparations at 45°C for 1 h had no significant effect on surface activity (Table 1). Decreasing the amount of SP-B to 0.5 weight % in 3 weight % SP-C(LKS) in DPPC:PG 7:3 (w/w) tended to increase γ_{\min} although the results did not reach statistical significance (Table 1). In contrast to SP-B, addition of 2 weight % KL₄ (Cochrane, C.G. and Revak, S.D. (1991) Science **254**, 566-568) to 3 weight % SP-C(LKS) in DPPC:PG:PA 68:22:9 (w/w) did not reduce γ_{\max} which remained relatively high at 41-42 mN/m.

Example 6

Comparison between mixtures containing dipalmitoylated and non-palmitoylated reference peptides

Surfactant preparations were prepared by adding 3% w/w SP-C(Leu) or dipalmitoylated SP-C(Leu) to each lipid mixtures, made of DPPC/PG/PA 68:22:9 w/w/w. The mixtures were evaporated under nitrogen and resuspended in 150 mmol/l NaCl at lipid concentrations of 10 mg/ml. In the samples in which a SP-B substituent was also used, 1% w/w of polymyxin B was added.

Mixtures containing dipalmitoylated SP-C(Leu), with or without polymyxin B, exhibit significant improvement especially in reducing γ_{\max} at 5 min and γ_{\min} at earlier time intervals.

Table 4: Surface properties

5 Surface tension of the mixtures was obtained with a pulsating bubble surfactometer. After two minutes of equilibration, the recordings were obtained at different periods, at 37° C, 50% surface compression and at a rate of 40 cycles per min.

Surfactant preparation			Surface tension (mN/m)					
SP-C(Leu)	Dipalm.	PxB	7,5 s		1 min		5 min	
SP-C(Leu)								
(weight %)	(weight %)	(weight %)	γ_{\min}	γ_{\max}	γ_{\min}	γ_{\max}	γ_{\min}	γ_{\max}
1			11	39	6,2	39	2	42
1		1	3	37	3	38	0	40
	1	-	1	34	1	35	1	36
	1	1	0	29	0	34	0	35

Example 7

In vivo determination

20 The effect of surfactant therapy on the mechanical properties of immature lungs was evaluated in 9 preterm newborn rabbits with a gestational age of 27 days. The animals were tracheotomized at birth and five of them received, via the tracheal cannula, twice 2.5 ml/kg of artificial surfactant containing DPPC, PG, and SP-C (LKS), with or without polymyxin B, in the proportions given above. Total phospholipid concentration of the

exogenous surfactant material was 40 mg/ml. Two animals serving as negative control received no material via the tracheal tube, and another two serving as positive control were treated with the same dose of modified natural surfactant (Curosurf, Chiesi Farmaceutici SpA, Parma, Italy), diluted to 40 mg/ml. One animal was treated with a mixture of DPPC and PG in saline (same concentrations as above) at a dose of 2.5 ml/kg. All animals were kept in body plethysmograph boxes at a temperature of 37°C and ventilated in parallel for 60 min with 100% oxygen, using a Servo Ventilator 900B (Siemens-Elema, Solna, Sweden) set at a frequency of 40 min and 50% inspiration time. Tidal volumes were measured with a pneumotachygraph connected to each plethysmograph box. The animals were ventilated with a standardized tidal volume of 8-10 ml/kg and without a positive end-expiratory pressure (PEEP). Lung-thorax compliance was defined as the ratio between tidal volume and peak inspiratory pressure, and expressed as ml/cm H₂O kg.

In comparison with the non-treated control animal, compliance was improved significantly in animals treated with the artificial surfactant, especially in the animal receiving surfactant containing polymyxin B. Notably, the improvement appears to be superior to that seen after treatment with a similar dose of modified natural surfactant (Fig. 3).

Brief description of figures.

Fig. 1. Amino acid sequences and helical wheel presentations of SP-C and its analogues.

The sequence of human SP-C is taken from Johansson, J. et al. (1988) FEBS Lett. **232**, 61-64 and that of SP-C(Leu) from

Nilsson, G., et al. (1998) Eur. J. Biochem. 255, 116-124). SP-C(LKS) is based on the primary structure of SP-C but all Val residues at the positions 16-28 with the exception of position 17 are replaced with Leu residues, Lys residues have been introduced at positions 17,22 and 27, and the palmitoylated Cys at positions 5 and 6 are replaced with Ser.

Fig. 2. Surface spreading of synthetic surfactant preparations.

Spreading kinetics of 3 weight % SP-C(LKS) (filled squares, solid line) and of 3 weight % SP-C(LKS) with addition of 2 weight % SP-B (open triangles, dotted line). All preparations were examined at a concentration of 10 mg /ml of DPPC/PG, 7:3 (w/w) in 150 mmol/l NaCl. The recordings were obtained with a Wilhelmy balance and each datapoint is the mean of three different recordings.

Fig. 3 In vivo results

Lung-thorax compliance in 5 premature newborn rabbits (gestational age of 27 days) ventilated with a standardized tidal volume of 8-10 ml/kg and without a positive end-expiratory pressure (PEEP). Compliance is improved significantly in treated animals. Addition of polymyxin B (PxB) appears to increase the effect of the artificial surfactant. The concentration of phospholipids is the same in all surfactant preparations, i.e. 40 mg/ml.

CLAIMS

1. SP-C analogues having general formula (I), according to one-letter amino acid code:



wherein:

X is an amino acid selected from the group consisting of V, I, L, Nle (norleucine);

10 B is an amino acid selected from the group consisting of ornithine, K, I, W, F, Y, Q, N;

Z is an amino acid selected from the group consisting of S, C, F where Ser or Cys residues are optionally linked via ester or thio-ester bonds with acyl group containing 12-22 carbon atoms.

a is an integer from 1 to 19;

15 b is an integer from 1 to 19;

c is an integer from 1 to 21;

d is an integer from 0 to 20;

e is 0 or 1;

f is 0 or 1;

20 n is 0 or 1;

m is 0 or 1,

with the following conditions:

- $n + m > 0$;

- $f \geq e$;

25 - $(\text{X}_a\text{B})(\text{X}_b\text{B})_n(\text{X}_c\text{B})_m\text{X}_d$ is a sequence having a maximum of 22 amino acids, preferably from 10 to 22 amino acids.

2. SP-C analogues according to claim 1, having formula (Ia):



3. SP-C analogues according to claim 1, having formula (Ib):



4. SP-C analogues according to claim 1, having formula (Ic)



5. SP-C analogues according to claim 1, having formula (Id)



6. SP-C analogues according to claim 1, having formula (Ie)



7. SP-C analogues according to claims 1-6, in which Ser residues
are acylated preferably with palmitoyl groups.

8. SP-C analogues according to claims 1-7, in which B is Lysine or
Phenylalanine and X is Leucine, Isoleucine or Norleucine.

9. SP-C analogues according to claim 8, selected from the
group consisting of:

SP-C (LKS) FGIPSSPVHLKRLILKLLLLKILLKLGALLMGL

SP-C (LKS)₁ FGIPSSPVHLKRLILKLLLLIKLLILGALLMGL

SP-C (LKS)₂ FGIPSSPVHLKRLILKLLLLLILLLILGALLMGL

SP-C (LKS)₃ FGIPSSPVHLKRLILKLLLLKLILLILGALLMGL

SP-C (LKS)₄ FGIPSSPVHLKRLILKLLLLLLIKLLILGALLMGL

SP-C (LFS) FGIPSSPVHLKRLILFLLLLFILLFLGALLMGL

10. A synthetic surfactant comprising at least one SP-C analogue
of formula (I) in admixture with lipids and phospholipids.

11. A synthetic surfactant according to claim 9, in which the
mixture lipids/phospholipids comprises DPPG, PG, PA.

12. A synthetic surfactant according to claims 10-11, further
comprising SP-B or an active derivative thereof or a polymyxin.

13. A synthetic surfactant according to claims 10-12, in form of
solution, dispersion, suspension, dry powder.

14. Use of SP-C analogues of claims 1-7 for the preparation of a synthetic surfactant to be used in all cases of surfactant deficiencies.

5 15. Use of a polymyxin, preferably polymyxin B for the preparation of an artificial surfactant according to claims 10-13, for the treatment of all cases of surfactant deficiencies or dysfunction, related pulmonary diseases such as pneumonia, bronchitis, asthma, meconium aspiration syndrome and also other diseases such as serous otitis media (glue ear).

10 16. Use according to claims 14 and 15, in which the surfactant deficiency is respiratory distress syndrome.

2/3

FIGURE 2





FIGURE 3



SEQUENCE LISTING

<110> Chiesi Farmaceutici spa

<120> Artificial peptides having surface activity and the use thereof in the preparation of artificial surfactant

<130> chiesi

<140>

<141>

<150> MI99A000275

<151> 1999-02-12

<160> 6

<170> PatentIn Ver. 2.1

<210> 1

<211> 35

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:syntetic peptides

<400> 1

Phe Gly Ile Pro Ser Ser Pro Val His Leu Lys Arg Leu Leu Ile Leu
1 5 10 15

Lys Leu Leu Leu Leu Lys Ile Leu Leu Leu Lys Leu Gly Ala Leu Leu
20 25 30

Met Gly Leu
35

<210> 2

<211> 35

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:syntetic peptides

<400> 2

Phe Gly Ile Pro Ser Ser Pro Val His Leu Lys Arg Leu Leu Ile Leu
1 5 10 15

Leu Lys Leu Leu Leu Leu Ile Lys Leu Leu Ile Leu Gly Ala Leu Leu
20 25 30

Met Gly Leu
35

<210> 3

<211> 35

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:syntetic
peptides

<400> 3

Phe Gly Ile Pro Ser Ser Pro Val His Leu Lys Arg Leu Leu Ile Leu
1 5 10 15

Lys Leu Leu Leu Leu Leu Ile Leu Leu Leu Ile Leu Gly Ala Leu Leu
20 25 30

Met Gly Leu
35

<210> 4

<211> 35

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:syntetic
peptides

<400> 4

Phe Gly Ile Pro Ser Ser Pro Val His Leu Lys Arg Leu Leu Ile Leu
1 5 10 15

Leu Leu Leu Leu Lys Leu Ile Leu Leu Leu Ile Leu Gly Ala Leu Leu
20 25 30

Met Gly Leu

35

<210> 5

<211> 35

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:syntetic
peptides

<400> 5

Phe Gly Ile Pro Ser Ser Pro Val His Leu Lys Arg Leu Leu Ile Leu
1 5 10 15

Leu Leu Leu Leu Leu Leu Ile Lys Leu Leu Ile Leu Gly Ala Leu Leu
20 25 30

Met Gly Leu
35

<210> 6

<211> 35

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:syntetic
peptides

<400> 6

Phe Gly Ile Pro Ser Ser Pro Val His Leu Lys Arg Leu Leu Ile Leu
1 5 10 15

Phe Leu Leu Leu Leu Phe Ile Leu Leu Leu Phe Leu Gly Ala Leu Leu
20 25 30

Met Gly Leu
35

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 00/01044

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C07K14/785 A61P11/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	PALMBLAD, MARIE ET AL.: "BIOPHYSICAL ACTIVITY OF AN ARTIFICIAL SURFACTANT CONTAINING AN ANALOGUE OF SURFACTANT PROTEIN (SP)-C AND NATIVE SP-B" BIOCHEM J (1999) 339(2) 381-386, April 1999 (1999-04), XP002139844 the whole document	1-16
X	WO 91 18015 A (CALIFORNIA BIOTECHNOLOGY INC) 28 November 1991 (1991-11-28)	1, 10-16
Y	claims 5, 8	7
X	EP 0 733 645 A (TOKYO TANABE CO) 25 September 1996 (1996-09-25)	1, 10-16
Y	abstract; claims	7
A	EP 0 368 823 A (KABIGEN AB) 16 May 1990 (1990-05-16)	
	-/-	



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

14 June 2000

Date of mailing of the international search report

29/06/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Cervigni, S

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 00/01044

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>TAKEI, TSUNETOMO ET AL.: "THE SURFACE PROPERTIES OF CHEMICALLY SYNTHESIZED PEPTIDES ANALOGOUS TO HUMAN PULMONARY SURFACTANT PROTEIN SP-C"</p> <p>BIOL PHARM BULL (1996) 19(10) 1247-1253, XP002139845</p> <p>-----</p>	

INTERNATIONAL SEARCH REPORT

...formation on patent family members

International Application No

PCT/EP 00/01044

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9118015 A	28-11-1991	US 5104853 A	14-04-1992
		CA 2083177 A	18-11-1991
		EP 0538273 A	28-04-1993
		JP 5509301 T	22-12-1993
		US 5385840 A	31-01-1995
EP 0733645 A	25-09-1996	AU 682738 B	16-10-1997
		AU 1199295 A	27-06-1995
		BG 100554 A	31-12-1996
		FI 962355 A	06-06-1996
		NO 962403 A	07-06-1996
		SK 71496 A	06-11-1996
		US 5827825 A	27-10-1998
		CA 2178345 A	15-06-1995
		CN 1136813 A	27-11-1996
		CZ 9601623 A	16-10-1996
		HU 74880 A,B	28-02-1997
		WO 9515980 A	15-06-1995
		NZ 277095 A	22-09-1997
		PL 314872 A	30-09-1996
EP 0368823 A	16-05-1990	AT 84799 T	15-02-1993
		AU 623180 B	07-05-1992
		AU 4299689 A	26-04-1990
		CA 2000893 A	18-04-1990
		DE 68904530 D	04-03-1993
		DE 68904530 T	08-07-1999
		JP 2145599 A	05-06-1990
		JP 2963923 B	18-10-1999
		SE 8803713 A	18-10-1988
		US 5455227 A	03-10-1995
		US 5223481 A	29-06-1993